

FIERCELY
PROTECTIVE



YOUR VET
HAS PRESCRIBED

NexGard[®]
COMBO

CONGRATULATIONS!

Your cat is now protected against these key parasites:

- ✓ FLEAS
- ✓ TICKS
- ✓ EAR MITES
- ✓ ROUNDWORMS
- ✓ TAPEWORMS
- ✓ LUNGWORMS
- ✓ NOTOEDRIC MANGE

So, let's take a quick look at why your vet prescribed NexGard® COMBO and how easy it is for you to maintain that protection.



NexGard® COMBO offers the broadest cat parasite protection available in a single treatment.

› Many cats don't like tablets so the simple-to-use applicator makes it easy to apply

› Can be used in cats and kittens in accordance with your vet's prescription

› Protects your cat inside and out



WHY

YOUR VET
HAS PRESCRIBED

NexGard®
COMBO

THESE PARASITES are a continuous health threat to your cat but can be controlled with NexGard® COMBO



THREAT FROM EAR MITES

- Ear mites are hard to spot! They are tiny animals and are a fairly common parasite in cats.
- Ear mites live on the skin of the ear canal and feed off ear wax and skin oils.
- Eggs take just four days to hatch and a further three weeks to develop into an adult mite ready to breed.

Ear mites are contagious to other cats and dogs in the household



YEAR-ROUND FLEAS

- Fleas are a common problem in cats and can carry diseases and tapeworms
- For every 5 fleas seen on your cat, there are 95 eggs, larvae or pupae hiding in your home or garden
- If your home is already contaminated your vet may recommend a household spray and treating any other pets with an appropriate product


Fleas can also bite humans and transmit disease¹



THREAT FROM TICKS

- Ticks are often so tiny they are almost impossible to find in your cat's fur
- Ticks can be found where cats like to explore – including gardens and urban areas
- Ticks feed on blood and can pass on diseases in the process²


Ticks can also transmit Lyme disease to people²



ROUNDWORM RISK

- Over 1 in 4 cats that go outdoors are infected and pass roundworm eggs in their poo³
- Roundworms can cause harm in both cats and people
- Children often play in areas where these roundworm eggs may be found. Roundworms can cause blindness and increase the risk of asthma and epilepsy⁴⁻⁶

Children and adults with a compromised immune system are especially at risk⁷



TAPEWORM INFESTATION

- Cats can be infected by swallowing fleas while grooming or by eating prey animals such as rodents
- Tapeworms live in cats' intestines and are made up of segments like flattened grains of rice connected together, each filled with eggs

Segments can break off and wiggle out through your cat's bottom. Yuk!

WHY

YOUR CAT AND YOUR FAMILY NEED

NexGard®
COMBO

HOW

TO APPLY


NexGard[®]
COMBO



- › Use scissors to cut the blister along the dotted line.



- › Pull the film away.



- › Pull back the plunger slightly.
- › Twist and pull off the cap.



- › Part the hair on the midline of the neck, between the base of the skull and the shoulder blades until the skin is visible and apply the entire contents directly onto the skin in one spot.



- › Wash hands.

Apply NexGard[®] COMBO where your cat cannot lick it off and in long haired breeds of cat special attention should be paid to apply NexGard[®] COMBO onto the skin and not on the fur.

Do not touch the treated area until no longer noticeable. It is recommended to treat animals in the evening to reduce contact and recently treated cats should not sleep with owners. For further information please read the package leaflet.



NexGard[®]
COMBO

NexGard[®] COMBO



References: 1. Wall, R. (2019) Flea and tick challenges - managing risk factors. *Vet Times*. 49(39). 2. Wright, I. (2012) Lyme disease: If you go down to the woods today are you in for a big surprise? *Companion Animal*. 17(6), 42-44. 3. Wright I, Stafford K, Coles G. The prevalence of intestinal nematodes in cats and dogs from Lancashire, north-west England. *J Small Anim Pract* 2016; 57: 393-395. 4. Ma G, Holland CV, Wang T, Hofmann A, Fan CK, Maizels RM, Hotez PJ, Gasser RB. Human toxocarasis. *Lancet Infect. Dis*. 2018; 18: e14-e24. 5. Quattrocchi G, Nicoletti A, Marin B, et al. Toxocarasis and epilepsy: systematic review and meta-analysis. *PLoS Negl Trop Dis* 2012;6:e1775. 6. Li L, Gao W, Yang X, et al. Asthma and toxocarasis. *Ann Allergy Asthma Immunol* 2014;113:187-192. 7. ESCCAP. *Worm Control in Dogs and Cats*. ESCCAP Guideline 01 Sixth Edition 2020.

NexGard[®] COMBO contains esafloxolaner, eprinomectin and praziquantel. UK: POM. Further information available in the SPC or from Boehringer Ingelheim Animal Health UK Ltd., RG12 8YS, UK. UK Tel: Pet 01344 746959 (sales) or 01344 746957 (technical). Email: vetenquiries@boehringer-ingelheim.com. NexGard[®] is a registered trademark of Boehringer Ingelheim Animal Health France SCS, used under licence. ©2024 Boehringer Ingelheim Animal Health UK Ltd. All rights reserved.
Date of preparation: Mar 2024. UI-FEL-0017-2024. Use Medicines Responsibly

